

Distance spectrum of graph compositions*

Gopalapillai Indulal

*Department of Mathematics, St. Aloysius College, Edathua
Alappuzha–689573, India*

Received 20 March 2009, accepted 15 May 2009, published online 19 June 2009

Abstract

The D -eigenvalues $\{\mu_1, \mu_2, \dots, \mu_p\}$ of a graph G are the eigenvalues of its distance matrix D and form the distance spectrum or D -spectrum of G denoted by $\text{spec}_D(G)$. In this paper we obtain the D -spectrum of the cartesian product of two distance regular graphs. The D -spectrum of the lexicographic product $G[H]$ of two graphs G and H when H is regular is also obtained. The D -eigenvalues of the Hamming graphs $\text{Ham}(d, n)$ of diameter d and order n^d and those of the C_4 nanotori, T_{k,m,C_4} are determined.

Keywords: Distance spectrum, Cartesian product, lexicographic product, Hamming graphs, C_4 nanotori.

Math. Subj. Class.: 05C12, 05C50

1 Introduction

Adjacency matrix of a graph and its spectrum have arisen as a natural tool with which one can study graphs and its structural properties. Also the adjacency spectrum find applications in quantum theory and chemistry [3]. The idea of distance matrix seems a natural generalization, with perhaps more specificity than that of an adjacency matrix. Distance matrix and their spectra have arisen independently from a data communication problem [7] studied by Graham and Pollack in 1971 in which the most important feature is the number of negative eigenvalues of the distance matrix. While the problem of computing the characteristic polynomial of adjacency matrix and its spectrum appears to be solved for many large graphs, the related distance polynomials have received much less attention. The distance matrix is more complex than the ordinary adjacency matrix of a graph since the distance matrix is a complete matrix (dense) while the adjacency matrix often is very sparse. Thus the computation of the characteristic polynomial of the distance matrix is

*This work was supported by the University Grants Commission of Government of India under the minor research project grant No: MRP(S)-399/08-09/KLMG019/UGC-SWRO.

E-mail address: indulalgopal@yahoo.com (Gopalapillai Indulal)

computationally a much more intense problem and, in general, there are no simple analytical solutions except for a few trees [6]. For this reason, distance polynomials of only trees have been studied extensively in the mathematical literature [6, 16].

The distance matrix of a graph has numerous applications to chemistry and other branches of science. The distance matrix, contains information on various walks and self-avoiding walks of chemical graphs, is immensely useful in the computation of topological indices such as the Wiener index, is useful in the computation of thermodynamic properties such as pressure and temperature coefficients and it contains more structural information compared to a simple adjacency matrix. In addition to such applications in chemical sciences, distance matrices find applications in music theory, ornithology, molecular biology, psychology, archeology etc. For a survey see [1] and also the papers cited therein.

Let G be a connected graph with vertex set $V(G) = \{u_1, u_2, \dots, u_p\}$. The distance matrix $D = D(G)$ of G is defined so that its (i, j) -entry is equal to $d_G(u_i, u_j)$, the distance (= length of the shortest path [2]) between the vertices u_i and u_j of G . The eigenvalues of $D(G)$ are said to be the D -eigenvalues of G and form the distance spectrum or the D -spectrum of G , denoted by $spec_D(G)$.

The characteristic polynomial of the D -matrix and the corresponding spectra have been considered in [4, 6, 7, 8]. For some recent works on D -spectrum see [9, 10, 11, 12, 13, 18].

For two graphs, the ordinary spectrum of graph compositions is well explored and generalized results of NEPS of graphs are presented in [3]. Such studies for the distance spectrum did not appear in literature yet and hence in this paper we present the following.

Let G and H be two graphs. Let $G + H$ and $G[H]$ denote the cartesian product and lexicographic product of G and H respectively [3].

In this paper we first derive the D -spectrum of $G + H$ and $G[H]$. By means of this, the distance spectrum of the Hamming graph and C_4 nanotori are obtained. A work of this type is reported here for the first time.

All graphs considered in this paper are simple and we follow [3] for spectral graph theoretic terminology and [2] for distance in graphs. The considerations in the subsequent sections are based on the applications of the following lemmas.

Lemma 1.1 ([3]). *Let G be an r -regular graph on p vertices with adjacency eigenvalues $r, \lambda_2, \dots, \lambda_p$. Then G and its complement \bar{G} have the same eigenvectors, and the eigenvalues of \bar{G} are $p - r - 1, -1 - \lambda_2, \dots, -1 - \lambda_p$.*

Lemma 1.2 ([5]). *The distance spectrum of the cycle C_n is given by*

n	greatest eigenvalue	j even	j odd
even	$\frac{n^2}{4}$	0	$-\operatorname{cosec}^2\left(\frac{\pi j}{n}\right)$
odd	$\frac{n^2 - 1}{4}$	$-\frac{1}{4} \sec^2\left(\frac{\pi j}{2n}\right)$	$-\frac{1}{4} \operatorname{cosec}^2\left(\frac{\pi j}{2n}\right)$

Definition 1.3 ([14]). The Hamming graph $Ham(d, n)$, $d \geq 2, n \geq 2$, of diameter d and characteristic n have vertex set consisting of all d -tuples of elements taken from an n -element set, with two vertices adjacent if and only if they differ in exactly one coordinate. $Ham(d, n)$ is equal to $\underbrace{K_n + K_n + \cdots + K_n}_d$, the cartesian product of K_n , the complete graph on n vertices, d times. $Ham(3, n)$ is referred to as a cubic lattice graph.

Lemma 1.4 ([17]). Let G and H be two connected graphs, and let $u = (u_1, u_2), v = (v_1, v_2) \in V(G) \times V(H)$. Let $G + H$ denote their cartesian product. Then

$$d_{G+H}(u, v) = d_G(u_1, v_1) + d_H(u_2, v_2).$$

2 The D -spectrum of $G + H$

In this section we derive the D -spectrum of the cartesian product of two distance regular graphs.

Theorem 2.1. Let G and H be two distance regular graphs on p and n vertices with distance regularity k and t respectively. Let $spec_D(G) = \{k, \mu_2, \mu_3, \dots, \mu_p\}$ and $spec_D(H) = \{t, \eta_2, \eta_3, \dots, \eta_n\}$. Then

$$spec_D(G + H) = \{nk + pt, n\mu_i, p\eta_j, 0\}$$

$i = 2, \dots, p, j = 2, \dots, n$ and 0 is with multiplicity $(p - 1)(n - 1)$.

Proof. Let D_G and D_H be the distance matrices of G and H respectively. Let $V(G) = \{u_1, u_2, \dots, u_p\}$ and $V(H) = \{v_1, v_2, \dots, v_n\}$. Then $D_G = [d_{ij}]$ and $D_H = [e_{ij}]$ where $d_{ij} = d_G(u_i, u_j)$ and $e_{ij} = d_H(v_i, v_j)$. Since G and H are distance regular graphs with distance regularities k and t respectively, we have

$$\sum_{j=1}^p d_{rj} = k \quad \text{and} \quad \sum_{j=1}^n e_{qj} = t \tag{2.1}$$

Also since G is distance regular, the all one column vector of order $p \times 1$ is the eigenvector corresponding to the greatest eigenvalue k of D_G . As D_G is real and symmetric, it is diagonalizable and hence admits an orthogonal basis B_G consisting of eigenvectors corresponding to its eigenvalues. Thus if μ_i is an eigenvalue of D_G which is different from k with an eigenvector $X_i = [x_i^1, x_i^2, \dots, x_i^p]^T \in B_G$, then $\sum_{j=1}^p x_i^j = 0$.

Let $u = (u_1, u_2), v = (v_1, v_2) \in V(G) \times V(H)$. Then by Lemma 1.4

$$d_{G+H}(u, v) = d_G(u_1, v_1) + d_H(u_2, v_2).$$

By a suitable ordering of vertices in $G + H$ and by virtue of Lemma 1.4, its D -matrix, C can be written in the form

$$\begin{aligned}
 C &= \begin{bmatrix} d_{11} + e_{11} & \cdots & d_{11} + e_{1n} & \cdots & \cdots & d_{1p} + e_{11} & \cdots & d_{1p} + e_{1n} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ d_{11} + e_{n1} & \cdots & d_{11} + e_{nn} & \cdots & \cdots & d_{1p} + e_{n1} & \cdots & d_{1p} + e_{nn} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ d_{p1} + e_{11} & \cdots & d_{p1} + e_{1n} & \cdots & \cdots & d_{pp} + e_{11} & \cdots & d_{pp} + e_{1n} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ d_{p1} + e_{n1} & \vdots & d_{p1} + e_{nn} & \cdots & \cdots & d_{pp} + e_{n1} & \cdots & d_{pp} + e_{nn} \end{bmatrix} \\
 &= \begin{bmatrix} d(u_1, u_1) \cdot J_n + D_H & d(u_1, u_2) \cdot J_n + D_H & \cdots & \cdots & d(u_1, u_p) \cdot J_n + D_H \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ d(u_p, u_1) \cdot J_n + D_H & d(u_p, u_2) \cdot J_n + D_H & \cdots & \cdots & d(u_p, u_p) \cdot J_n + D_H \end{bmatrix} \\
 &= D_G \otimes J_n + J_p \otimes D_H
 \end{aligned}$$

where \otimes denotes the tensor product of matrices.

Now we find the eigenvalues of C by considering eigenvectors associated with them. The following relation for matrices is well known [15]. For the matrices A, B, C and D

$$(A \otimes B) \cdot (C \otimes D) = (AC) \otimes (BD)$$

whenever the products AC and BD exist.

Let $\mathbf{1}_G$ denote the all one eigenvector corresponding to the eigenvalue k of G and $\mathbf{1}_H$ the all one eigenvector corresponding to the eigenvalue t of H . Then

$$D_G \cdot \mathbf{1}_G = k\mathbf{1}_G \quad \text{and} \quad D_H \cdot \mathbf{1}_H = t\mathbf{1}_H$$

Therefore

$$\begin{aligned}
 C \cdot (\mathbf{1}_G \otimes \mathbf{1}_H) &= (D_G \otimes J_n + J_p \otimes D_H) \cdot (\mathbf{1}_G \otimes \mathbf{1}_H) \\
 &= (D_G \cdot \mathbf{1}_G) \otimes (J_n \mathbf{1}_H) + (J_p \mathbf{1}_G) \otimes (D_H \cdot \mathbf{1}_H) \\
 &= k\mathbf{1}_G \otimes n\mathbf{1}_H + p\mathbf{1}_G \otimes t\mathbf{1}_H \\
 &= nk \cdot (\mathbf{1}_G \otimes \mathbf{1}_H) + pt \cdot (\mathbf{1}_G \otimes \mathbf{1}_H) \\
 &= (nk + pt) \cdot (\mathbf{1}_G \otimes \mathbf{1}_H)
 \end{aligned}$$

showing that $\mathbf{1}_G \otimes \mathbf{1}_H$ is the eigenvector corresponding to the eigenvalue $nk + pt$ of C .

Let X_i be the eigenvector corresponding to the eigenvalue μ_i of D_G . Then $X_i \otimes \mathbf{1}_H$ is the eigenvector corresponding to the eigenvalue $n\mu_i$ of C . For

$$\begin{aligned}
 C \cdot (X_i \otimes \mathbf{1}_H) &= (D_G \otimes J_n + J_p \otimes D_H) \cdot (X_i \otimes \mathbf{1}_H) \\
 &= (D_G \cdot X_i) \otimes (J_n \mathbf{1}_H) + (J_p X_i) \otimes (D_H \cdot \mathbf{1}_H) \\
 &= \mu_i X_i \otimes n\mathbf{1}_H + 0 \otimes t\mathbf{1}_H \\
 &= n\mu_i (X_i \otimes \mathbf{1}_H)
 \end{aligned}$$

Similarly if Z_j is an eigenvector corresponding to the eigenvalue η_j of D_H , then $\mathbf{1}_G \otimes Z_j$ is an eigenvector corresponding to the eigenvalue $p\eta_j$ of C .

In addition to these eigenvalues we can see that 0 appears to be an eigenvalue with multiplicity $(p-1)(n-1)$. For let R_p^i , $i = 2, 3, \dots, p$ be the $(p-1)$ linearly independent eigenvectors corresponding to the eigenvalue 0 of J_p and T_n^j , $j = 2, 3, \dots, n-1$ be the $(n-1)$ linearly independent eigenvectors corresponding to the eigenvalue 0 of J_n . Then the $(p-1)(n-1)$ vectors $R_p^i \otimes T_n^j$ are linearly independent and are the eigenvectors corresponding to 0 of C . For

$$\begin{aligned} C \cdot (R_p^i \otimes T_n^j) &= (D_G \otimes J_n + J_p \otimes D_H) \cdot (R_p^i \otimes T_n^j) \\ &= (D_G \cdot R_p^i) \otimes (J_n \cdot T_n^j) + (J_p R_p^i) \otimes (D_H \cdot T_n^j) \\ &= (D_G \cdot R_p^i) \otimes 0 + 0 \otimes (D_H \cdot T_n^j) \\ &= 0 \end{aligned}$$

Now the pn vectors $X_i \otimes \mathbf{1}_H$, $\mathbf{1}_G \otimes Z_j$ and $R_p^i \otimes T_n^j$ are linearly independent and as C has a basis consisting of linearly independent eigenvectors, the theorem follows. \square

2.1 The D -spectrum of $Ham(d, n)$

In [14], the ordinary spectrum of the cubic lattice graph is obtained. In this section we use Theorem 2.1 to obtain the D -spectrum of $Ham(d, n)$.

Theorem 2.2. *Let $Ham(d, n)$ be the Hamming graph of characteristic n . Then the D -eigenvalues of $Ham(d, n)$ are $dn^{d-1}(n-1)$, 0 and $-n^{d-1}$ with multiplicities 1, $n^d - d(n-1) - 1$ and $d(n-1)$ respectively.*

Proof. The graph K_n is distance regular with distance regularity $n-1$. Now the proof follows by repeated application of Theorem 2.1 and from the ordinary spectrum of K_n [3]. \square

2.2 The D -spectrum of the C_4 nanotori, T_{k,m,C_4}

The graph $C_k + C_m$ where both k and m are odd is defined as the C_4 nanotori, T_{k,m,C_4} .

Theorem 2.3. *The distance spectrum of the C_4 nanotori, T_{k,m,C_4} consists of the following numbers*

$$\begin{aligned} &\frac{(m+k)(mk-1)}{4}, -\frac{m}{4} \sec^2\left(\frac{\pi j}{2k}\right), -\frac{m}{4} \operatorname{cosec}^2\left(\frac{\pi r}{2k}\right), \\ &-\frac{k}{4} \sec^2\left(\frac{\pi t}{2m}\right), -\frac{k}{4} \operatorname{cosec}^2\left(\frac{\pi l}{2m}\right) \end{aligned}$$

where $j \in \{1, 2, \dots, k-1\}$ and even, $r \in \{1, 2, \dots, k-1\}$ and odd $t \in \{1, 2, \dots, m-1\}$ and even and $l \in \{1, 2, \dots, m-1\}$ and odd together with 0 of multiplicity $(m-1)(k-1)$.

Proof. The cycle C_{2n+1} is distance regular with distance regularity $n(n+1)$. Now the proof follows from Theorem 2.1 and Lemma 1.2. \square

3 The D -spectrum of $G[H]$

In this section we obtain the distance spectrum of the lexicographic product $G[H]$ of two graphs G and H . The following definition of the lexicographic product of G and H is from [3].

Definition 3.1. Let G and H be two graphs on vertex sets $V(G) = \{u_1, u_2, \dots, u_p\}$ and $V(H) = \{v_1, v_2, \dots, v_n\}$ respectively. Then their lexicographic product $G[H]$ is a graph defined by $V(G[H]) = V(G) \times V(H)$, the cartesian product of $V(G)$ and $V(H)$ in which $u = (u_1, v_1)$ be adjacent to $v = (u_2, v_2)$ if and only if either

1. u_1 be adjacent to v_1 in G or
2. $u_1 = v_1$ and u_2 be adjacent to v_2 in G .

Distance in $G[H]$

We prove the following lemma on distance in lexicographic product of graphs.

Lemma 3.2. *Let G and H be two connected graphs with atleast two vertices and let $u = (u_1, v_1), v = (u_2, v_2) \in V(G) \times V(H)$. Then*

$$d_{G[H]}(u, v) = \begin{cases} d_G(u_1, u_2) & \text{if } u_1 \neq u_2 \\ 1 & \text{if } u_1 = u_2 \text{ and } v_1 \text{ adjacent to } v_2 \\ 2 & \text{if } u_1 = u_2 \text{ and } v_1 \text{ not adjacent to } v_2 \end{cases}$$

Proof. We show that in the corresponding composition there exist a path between u and v of length as given in the lemma. Let $d_G(u_1, u_2) = t$ and $u_1 = s_0, s_1, \dots, s_t = u_2$ be the shortest $u_1 - u_2$ path in G .

Let $u = (u_1, v_1), v = (u_2, v_2) \in V(G) \times V(H)$ and $u_1 \neq u_2$. Since the successive ordered pairs in any $u - v$ path can change both the coordinates and also as u_2 is reachable from u_1 by not less than t steps, any $u - v$ path in $G[H]$ is of length atleast t .

Now the following $u - v$ path in $G[H]$ is of length t .

$P : u = (s_0, v_1), (s_1, v_2), (s_2, v_2), \dots, (s_t, v_2) = v$. Thus $d_{G[H]}(u, v) = d_G(u_1, u_2)$ if $u_1 \neq u_2$.

Now suppose $u_1 = u_2$ and v_1 be adjacent to v_2 . Then by the definition of $G[H]$, we have $d_{G[H]}(u, v) = 1$.

Now suppose $u_1 = u_2$ and v_1 is not adjacent to v_2 . Let s_1 be adjacent to u_1 in G . Then u is not adjacent to v and $u = (u_1, v_1), (s_1, v_2), (u_1, v_2) = v$ is a $u - v$ path of length 2. Thus $d_{G[H]}(u, v) = 2$. Hence the Lemma. \square

Theorem 3.3. *Let G be a graph with D -matrix D_G and H , an r -regular graph with an adjacency matrix A . Let $spec_D(G) = \{\mu_1, \mu_2, \dots, \mu_p\}$ and the ordinary spectrum of H be $\{r, \lambda_2, \lambda_3, \dots, \lambda_n\}$. Then*

$$spec_D G[H] = \left(\begin{array}{cc} n\mu_i + 2n - r - 2 & -(\lambda_j + 2) \\ 1 & p \end{array} \right), \quad i = 1 \text{ to } p \text{ and } j = 2 \text{ to } n - 1$$

Proof. Using Lemma 3.2 and by a suitable ordering of vertices of $G[H]$, its D -matrix F , can be written in the form

$$F = \begin{bmatrix}
 & & & d_{12} & \cdots & d_{12} & d_{13} & \cdots & d_{13} & \cdots & \cdots & d_{1p} \\
 & A + 2\bar{A} & & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
 & & & d_{12} & d_{12} & d_{12} & d_{13} & \cdots & d_{13} & \cdots & \cdots & d_{1p} \\
 d_{21} & \cdots & d_{21} & & & & \cdots & \cdots & \cdots & d_{2p} & \cdots & d_{2p} \\
 \vdots & \vdots & \vdots & & & & & & & \vdots & \vdots & \vdots \\
 d_{21} & \cdots & d_{21} & & A + 2\bar{A} & & & & & d_{2p} & \cdots & d_{2p} \\
 \vdots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
 & & & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
 d_{p1} & \cdots & d_{p1} & \cdots & \cdots & \cdots & \ddots & \cdots & \cdots & & & \\
 \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
 d_{p1} & \cdots & d_{p1} & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & A + 2\bar{A} & \\
 \end{bmatrix} \\
 = D_G \otimes J_n + I_p \otimes (A + 2\bar{A})$$

where \bar{A} denote the adjacency matrix of \bar{G} .

Since H is r -regular, the all one column vector $\mathbf{1}$ of order $n \times 1$ is an eigenvector of A with an eigenvalue r . Then by Lemma 1.1, the all one vector $\mathbf{1}$ is an eigenvector of $A + 2\bar{A}$ with an eigenvalue $2n - r - 2$. Similarly if λ_j is any other eigenvalue of A with eigenvector Y_j , then Y_j is an eigenvector of $A + 2\bar{A}$ with eigenvalue $-(\lambda_j + 2)$ and that Y_j is orthogonal to $\mathbf{1}$.

Let $X_i = [x_1^i \ x_2^i \ \dots \ x_p^i]^T$ be an eigenvector corresponding to the eigenvalue μ_i of D_G . Therefore

$$D_G \cdot X_i = \mu_i X_i$$

Now

$$\begin{aligned}
 F \cdot (X_i \otimes \mathbf{1}_n) &= (D_G \otimes J_n + I_p \otimes (A + 2\bar{A})) (X_i \otimes \mathbf{1}_n) \\
 &= (D_G \cdot X_i) \otimes (J_n \cdot \mathbf{1}_n) + (I_p \cdot X_i) \otimes (A + 2\bar{A}) \cdot \mathbf{1}_n \\
 &= \mu_i X_i \otimes n\mathbf{1}_n + X_i \otimes (2n - r - 2) \mathbf{1}_n \\
 &= n\mu_i (X_i \otimes \mathbf{1}_n) + (2n - r - 2) (X_i \otimes \mathbf{1}_n) \\
 &= (n\mu_i + 2n - r - 2) (X_i \otimes \mathbf{1}_n)
 \end{aligned}$$

Therefore $n\mu_i + 2n - r - 2$ is an eigenvalue of F with eigenvector $X_i \otimes \mathbf{1}_n$. As Y_j is orthogonal to $\mathbf{1}$, we have $J_n \cdot Y_j = 0$ for each $j = 2, 3, \dots, n$.

Let $\{Z_k\}, k = 1, 2, \dots, p$ be the family of p linearly independent eigenvectors associated with the eigenvalue 1 of I_p . Then for each $j = 2, 3, \dots, n$, the p vectors $Z_k \otimes Y_j$ are eigenvectors of F with eigenvalue $-(\lambda_j + 2)$. For

$$\begin{aligned}
 F \cdot (Z_k \otimes Y_j) &= (D_G \otimes J_n + I_p \otimes (A + 2\bar{A})) (Z_k \otimes Y_j) \\
 &= (D_G \cdot Z_k) \otimes (J_n \cdot Y_j) + (I_p \cdot Z_k) \otimes (A + 2\bar{A}) \cdot Y_j \\
 &= 0 + Z_k \otimes -(\lambda_j + 2) Y_j \\
 &= -(\lambda_j + 2) \cdot (Z_k \otimes Y_j)
 \end{aligned}$$

Also the pn vectors $X_i \otimes \mathbf{1}_n$ and $Z_k \otimes Y_j$ are linearly independent. As the eigenvectors belonging to different eigenvalues are linearly independent and as F has a basis consisting entirely of eigenvectors, the theorem follows. \square

Acknowledgements

The author is indebted to the anonymous referees for their valuable comments and suggestions which led to an improved presentation of the results.

References

- [1] K. Balasubramanian, Computer Generation of Distance Polynomials of Graphs, *Journal of Computational Chemistry* **11** (1990), 829–836.
- [2] F. Buckley and F. Harary, *Distance in Graphs*, Addison Wesley, 1990.
- [3] D. M. Cvetković, M. Doob and H. Sachs, *Spectra of Graphs. Theory and Applications*, Academic Press, New York – London, 1980.
- [4] M. Edelberg, M. R. Garey and R. L. Graham, On the distance matrix of a tree, *Discrete Math.* **14** (1976), 230–39.
- [5] P. W. Fowler, G. Caporossi and P. Hansen, Distance matrices, Wiener indices, and related invariants of fullerenes, *J. Phys. Chem. A.* **105** (2001), 6232–6242.
- [6] R. L. Graham and L. Lovasz, Distance matrix polynomials of trees, *Adv. Math.* **29** (1978), 60–88.
- [7] R. L. Graham and H. O. Pollak, On the addressing problem for loop switching, *Bell System Tech. J.* **50** (1971), 2495–2519.
- [8] R. L. Graham and H. O. Pollak, On embedding graphs in squashed cubes, in: Y. Alavi, D. R. Lick and A. T. White (eds.), *Graph theory and applications* (Lecture notes in Mathematics, vol. 303), Springer, Berlin, 1972, 99–110.
- [9] G. Indulal, I. Gutman and A. Vijayakumar, On distance energy of graphs, *MATCH Commun. Math. Comput. Chem.* **60** (2008), 461–472.
- [10] G. Indulal and I. Gutman, On the distance spectra of some graphs, *Math. Commun.* **13** (2008), 123–131.
- [11] G. Indulal and I. Gutman, D -equienergetic self-complementary graphs, *Kragujevac J. Math.*, in press.
- [12] G. Indulal, D -spectra and D -energy of the complements of iterated line graphs of regular graphs, communicated.
- [13] G. Indulal, Sharp bounds on the distance spectral radius and the distance energy of graphs, *Lin. Algebra Appl.* **430** (2009), 106–113.
- [14] R. Laskar, Eigenvalues of the adjacency matrix of cubic lattice graphs, *Pacific J. Math.* **29** (1969), 623–629.
- [15] M. Marcus and H. Mine, *A survey of Matrix theory and Matrix inequalities*, Allyn and Bacon, Inc., Boston, Mass., 1964.
- [16] B. D. McKay, On the spectral characterization of trees, *Ars Combin.* **3** (1977), 219–232.
- [17] D. Stevanović, Distance Regularity of Compositions of Graphs, *Appl. Math. Lett.* **17** (2004), 337–343.
- [18] D. Stevanović and G. Indulal, The distance spectrum and energy of the compositions of regular graphs, *Appl. Math. Lett.* **22** (2009), 1136–1140.