

SOME NEW INTEGRAL GRAPHS

G.Indulal, A.Vijayakumar

The eigenvalue of a graph is the eigenvalue of its adjacency matrix. A graph G is integral if all of its eigenvalues are integers. In this paper some new classes of integral graphs are constructed.

1. INTRODUCTION

Let G be a graph with $|V(G)| = n$ and adjacency matrix A . The eigenvalues of A are called the eigenvalues of G and form the spectrum of G denoted by $spec(G)$ in CVETKOVIĆ [2]. The graph G is integral if $spec(G)$ consists of only integers.

In BALIŃSKA [1] constructions and properties of integral graphs are discussed in detail. The graphs K_p and $K_{p,p}$ are examples of integral graphs. Some recent work on these lines pertaining to the class of trees is found in WANG [4]. Moreover, several graph operations such as Cartesian product, Strong sum and Product on integral graphs can be used for constructing infinite families of integral graphs, BALIŃSKA [1].

In this paper we provide some new constructions to obtain integral graphs. All graph theoretic terminology is from CVETKOVIĆ [2].

2. MAIN THEOREMS

The characteristic polynomial of G , $|\lambda I - A|$ is denoted by $P(G)$. A graph G , is said to be rooted at u if u is a specified vertex of G . We use the following lemmas in this paper.

Lemma 1 (SCHWENK [3]). *Let G and H be graphs rooted at u and v respectively.*

1. *Let F be the graph obtained by making u and v adjacent. Then*

$$P(F) = P(G)P(H) - P(G - u)P(H - v).$$

2000 Mathematics Subject Classification. 05C50.

Key Words and Phrases. Adjacency matrix, eigenvalue, integral graphs.

2. Let F' be the graph obtained by identifying u and v . Then

$$P(F') = P(G)P(H - v) + P(G - u)P(H) - xP(G - u)P(H - v).$$

Lemma 2 (CVETKOVIĆ [2]). Let M, N, P and Q be matrices with M invertible. Let $S = \begin{bmatrix} M & N \\ P & Q \end{bmatrix}$. Then $\det S = |M| |Q - PM^{-1}N|$.

Lemma 3 (CVETKOVIĆ [2]). Let G be an r -regular connected graph on p vertices with $r = \lambda_1, \lambda_2, \dots, \lambda_m$ as the distinct eigenvalues. Then there exists a polynomial $Q(x)$ such that $Q\{A(G)\} = J$, where J is the all one square matrix of order p and $Q(x)$ is given by $Q(x) = p \times \frac{(x - \lambda_2)(x - \lambda_3) \cdots (x - \lambda_m)}{(r - \lambda_2)(r - \lambda_3) \cdots (r - \lambda_m)}$, so that $Q(r) = p$ and $Q(\lambda_i) = 0$, for all $\lambda_i \neq r$.

Definition 1 (CVETKOVIĆ [2]). Let G be an r_1 -regular graph on p_1 vertices and H , an r_2 -regular graph on p_2 vertices. Then the complete product of G and H , denoted by $G \nabla H$ is obtained by joining every vertex of G to every vertex of H .

Note 1. The characteristic polynomial of $G \nabla H$ is given by

$$P(G \nabla H) = \frac{P(G)P(H)}{(x - r_1)(x - r_2)} (x^2 - (r_1 + r_2)x + r_1r_2 - p_1p_2).$$

Notation 1. Let $k *_G H$ denote the graph obtained by joining roots in k copies of H to all vertices of G . This graph can be obtained by first forming the complete product $G \nabla \overline{K_k}$ and then successively identifying the vertices in $\overline{K_k}$ one by one with roots in the k copies of H .

Let F_k^t , $t \leq k$ denote the graph obtained by identifying roots of t copies of H with t vertices of $\overline{K_k}$ in $G \nabla \overline{K_k}$. Then $F_k^0 = G \nabla \overline{K_k}$ and $F_k^k = k *_G H$.

Notation 2. $H_k = k \bullet H$, denote the graph obtained by identifying the root v in k copies of H .




Figure 1. F_3^2 when $G = K_{1,2}$ and $H = K_3$




Figure 2. H_4 when $H = K_3$

Theorem 1. Let G be an m -regular graph on p vertices and H be rooted at v . Then with the notations as described above

$$P(F_k^t) = \frac{P(G)}{(x - m)} x^{k-(t+1)} (P(H))^{t-1} (P(H)(x(x - m) - p(k - t)) - tpxP(H - v)).$$

Proof. We shall prove the Theorem by mathematical induction on t .

When $t = 0$, $F_k^0 = G\nabla \overline{K_k}$ and in this case

$$P(F_k^0) = \frac{P(G)}{(x-m)} x^{k-1} (x(x-m) - pk),$$

which is true from Note 1.

Now assume that the Theorem is true when $t = r < k$. Thus

$$P(F_k^r) = \frac{P(G)}{(x-m)} x^{k-(r+1)} (P(H))^{r-1} \left(P(H)(x(x-m) - p(k-r)) - rpxP(H-v) \right).$$

Now F_k^{r+1} is the graph obtained from F_k^r by identifying the $(r+1)^{th}$ vertex of $\overline{K_k}$ in $G\nabla \overline{K_k}$ with the root v in the $(r+1)^{th}$ copy of H . Now by Lemma 1 and by the induction hypothesis

$$\begin{aligned} P(F_k^{r+1}) &= P(F_k^r)P(H-v) + P(F_{k-1}^r)P(H) - xP(F_{k-1}^r)P(H-v) \\ &= \frac{P(G)}{(x-m)} x^{k-(r+1)} (P(H))^{r-1} \left(P(H)(x(x-m) - p(k-r)) - rpxP(H-v) \right) P(H-v) \\ &\quad + \frac{P(G)}{(x-m)} x^{k-1-(r+1)} (P(H))^{r-1} \left(P(H)(x(x-m) - p(k-1-r)) - rpxP(H-v) \right) P(H) \\ &\quad - xP(H-v) \frac{P(G)}{(x-m)} x^{k-(r+2)} (P(H))^{r-1} \left(P(H)(x(x-m) - p(k-1-r)) - rpxP(H-v) \right) \\ &= \frac{P(G)}{(x-m)} x^{k-(r+2)} (P(H))^r \left(P(H)(x(x-m) - p(k-(r+1))) - (r+1)pxP(H-v) \right). \end{aligned}$$

Thus the Theorem is true for $t = r + 1$. Hence by mathematical induction the Theorem follows. \square

Corollary 1.

$$P(k *_G H) = P(F_k^k) = \frac{P(G)}{(x-m)} (P(H))^{k-1} (P(H)(x-m) - pkP(H-v)).$$

Theorem 2. *The characteristic polynomial of H_k is given by*

$$P(H_k) = (P(H-v))^{k-1} (kP(H) - (k-1)xP(H-v)).$$

Proof. We shall prove the Theorem by mathematical induction on k and by Lemma 1. The Theorem is trivially true when $k = 1$. Assume that the result is true for $t < k$. Thus

$$P(H_t) = (P(H - v))^{t-1} (tP(H) - (t - 1)xP(H - v)).$$




Figure 3.

Now

$$\begin{aligned} P(H_{t+1}) &= P(H)(P(H - v))^t + P(H - v)P(H_t) - xP(H - v)(P(H - v))^t \\ &= P(H)(P(H - v))^t + P(H - v) \left((P(H - v))^{t-1} (tP(H) \right. \\ &\quad \left. - (t - 1)xP(H - v)) \right) - x(P(H - v))^{t+1} \\ &= (P(H - v))^t ((t + 1)P(H) - txP(H - v)). \end{aligned}$$

Hence the Theorem is true for $t+1$ and by mathematical induction Theorem follows. \square

3. SOME NEW INTEGRAL GRAPHS

In this section we shall give some new constructions of integral graphs.

Construction 1. Let G be any m - regular integral graph and H be K_{m+2} . Then by Theorem 1, the graph $k *_G H$ is integral if and only if the roots of $(x - m - 1)(x + 1) - pk = 0$ are integers. That happens if and only if $(m + 2)^2 + 4pk$ is a perfect square. Thus for $k = \frac{h^2 - (m + 2)^2}{4p}$, $h > m + 2$, we get an infinite family of integral graphs.

EXAMPLE 1.




Figure 4. $G = K_3$, $m = 2$,
 $H = K_4$, $k = 4$




Figure 5. $G = C_4$, $m = 2$,
 $H = K_4$, $k = 3$

Construction 2. Let $G = K_{m,n}$ with any vertex in the n vertex set as root. Then by Theorem 2, G_k is integral if and only if both $m(n-1)$ and $m(n-1) + mk$ are perfect squares. Now $m = t$; $n = t + 1$; $k = 3t$ is a feasible solution.

Construction 3. Let $G = K_4 - e$ rooted at any of the two non adjacent vertices. Then by Theorem 2, G_k is integral if and only if $8k + 9$ is a perfect square. Then for integer k of the form $k = \frac{t^2 - 9}{8}$, $t \geq 4$, we get an infinite family of integral graphs.

EXAMPLE 2.




Figure 6. $G = K_{2,3}$, $k = 6$
 $H = K_4$, $k = 4$

EXAMPLE 3.




Figure 7. $G = K_4 - e$, $k = 5$
 $H = K_4$, $k = 3$

4. SOME OPERATIONS ON GRAPHS

In this section we define some operations on a regular graph and thus provide some infinite families of integral graphs. Let G be a connected r -regular graph on p vertices and q edges whose adjacency matrix is A and spectrum $\{\lambda_1 = r, \lambda_2, \lambda_3, \dots, \lambda_p\}$.

Operation 1. Corresponding to each edge of G introduce a vertex and make it adjacent to vertices incident with it. Now introduce k isolated vertices and make all of them adjacent to vertices of G only.

Operation 2. Form the subdivision graph of G . Introduce k vertices and make all of them adjacent to vertices of G only.

Operation 3. Form the subdivision graph of G and add a pendant edge at each vertex of G . Introduce k vertices and make all of them adjacent to vertices of G only.

Theorem 3. Let G be a connected r -regular graph on p vertices and q edges with adjacency matrix A and spectrum $\{r, \lambda_2, \lambda_3, \dots, \lambda_p\}$. Let F_i be the graph obtained from G by operation i , $i = 1$ to 3. Then

$$\begin{aligned} \text{spec}(F_1) &= \begin{pmatrix} 0 & \frac{r \pm \sqrt{r^2 + 4(pk + 2r)}}{2} & \frac{\lambda_2 \pm \sqrt{\lambda_2^2 + 4(\lambda_2 + r)}}{2} & \dots & \frac{\lambda_p \pm \sqrt{\lambda_p^2 + 4(\lambda_p + r)}}{2} \\ k + q - p & \text{each once} & \text{each once} & \dots & \text{each once} \end{pmatrix} \\ \text{spec}(F_2) &= \begin{pmatrix} 0 & \pm\sqrt{(pk + 2r)} & \pm\sqrt{(\lambda_2 + r)} & \dots & \pm\sqrt{(\lambda_p + r)} \\ k + q - p & \text{each once} & \text{each once} & \dots & \text{each once} \end{pmatrix} \\ \text{spec}(F_3) &= \begin{pmatrix} 0 & \pm\sqrt{(pk + 2r + 1)} & \pm\sqrt{(\lambda_2 + r + 1)} & \dots & \pm\sqrt{(\lambda_p + r + 1)} \\ k + q & \text{each once} & \text{each once} & \dots & \text{each once} \end{pmatrix} \end{aligned}$$

Proof. The proof follows from the Table 1 which gives the adjacency matrix and characteristic polynomial of F_i , $i = 1, 2, 3$.

Table 1

Graph	Adjacency matrix	Characteristic polynomial
F_1	$\begin{bmatrix} A & R & J_{p \times k} \\ R^T & 0 & 0 \\ J_{k \times p} & 0 & 0 \end{bmatrix}$	$x^{q+k-p} \prod_{i=1}^p (x^2 - \lambda_i x - (kJ + \lambda_i + r))$
F_2	$\begin{bmatrix} 0 & R & J_{p \times k} \\ R^T & 0 & 0 \\ J_{k \times p} & 0 & 0 \end{bmatrix}$	$x^{q+k-p} \prod_{i=1}^p (x^2 - (kJ + \lambda_i + r))$
F_3	$\begin{bmatrix} 0 & R & I & J_{p \times k} \\ R^T & 0 & 0 & 0 \\ I & 0 & 0 & 0 \\ J_{k \times p} & 0 & 0 & 0 \end{bmatrix}$	$x^{q+k} \prod_{i=1}^p (x^2 - (kJ + \lambda_i + r + 1))$

where R is the incidence matrix with $RR^T = A + rI$ and J is the the all one matrix as in Lemma 3. □

EXAMPLES:

- $G = K_{p,p}$. F_1 is integral if and only if $p = t^2$, and $k = 2\ell^2 \pm \ell t - 1$, $\ell \geq t$, $t \geq 1$.

2. $G = K_{p,p}$. F_2 is integral if and only if $p = t^2$, and $k = 2h^2 - 1$, $t \geq 1$, $h \geq 1$.
3. $G = K_p$. F_3 is integral when $p = t^2$, and $k = t^2h^2 \pm 2h - 2$, $t \geq 1$, $h \geq 1$.
4. $G = K_{p,p}$. F_3 is integral when $p = t^2 - 1$, and $k = 2(t^2 - 1)h^2 \pm 2h - 1$, $t \geq 1$, $h \geq 1$.

Acknowledgments. We thank the referee for some suggestions. The first author thanks the University Grants Commission (India) for providing fellowship under the Faculty Improvement Programme.

REFERENCES

1. K. BALIŃSKA, D. CVETKOVIĆ, Z. RADOSAVLJEVIĆ, S. SIMIĆ, D. STEVANOVIĆ: *A Survey On Integral Graphs*. Univ. Beograd. Publ. Elektrotehn. Fak., Ser. Mat., **13** (2002), 42-65.
2. D. M. CVETKOVIĆ, M. DOOB, H. SACHS: *Spectra of Graphs-Theory and Applications*. Academic Press, New York, 1980.
3. A. J. SCHWENK: *On the eigenvalue of a graph*. In: L. W. BEINEKE, R. J. WILSON (Eds.): *Selected Topics in Graph Theory*. Academic Press, London, 1978, 307- 336.
4. L. WANG: *Integral Graphs and Integral Trees*. Ph.D.Thesis. University of Twente, 2005.

Department of Mathematics,
Cochin University of Science and Technology
Cochin-682 022, India.

E-mails: indulalgopal@cusat.ac.in, vijay@cusat.ac.in

(Received October 11, 2006)